
# **Ecotox Report for Case # P-19-0139**

#### General



### **Fate Summary Statement**

```
Fate Summary P 19 0139

Statement: FATE: PMN Substance: with MP < □ °C (E)

S

Hydrolysis Half-life ≤ min

VP torr at 25 °C (NOMO5)

BP = □ °C (M)

POTW removal (%) PMN 90 99 via hydrolysis; then Hyd Pdt Perfluoro acid 0;

Hyd Pdt 60 via sorption

Time for complete ultimate aerobic biodeg = Hyd Pdt Perfluoro acid > mo;

Hyd Pdt mo

Sorption to soils/sediments = Hyd Pdt Perfluoro acid low; Hyd Pdt moderate
```

PBT Potential: PMN P1B1; Hyd Pdt Perfluoro acid P3BU; Hyd Pdt P3B1 FATE: Migration to ground water = PMN negl; Hyd Pdt Perfluoro acid rapid: Hyd Pdt moderate Analogue found (include identifier and database) Relevant Structure(s) Water Perlfuoro product and Landfill Perlfuoro product and Air / Incineration Parent Parent % incineration 99.9% (hazardous waste incinerator) Environmental Fate Determination PMN #: P-19-0139 Summary: EPA estimated that the new chemical substance could have limited persistence and a low potential for bioaccumulation, such that repeated exposures are not expected to cause food-chain effects via accumulation in exposed organisms. The hydrolysis product (perfluoro acid) is estimated to be very persistent in the environment and its bioaccumulation potential is unknown; therefore, it is not known whether repeated exposures to it will cause food chain effects via accumulation in exposed organisms. Although EPA estimated that the hydrolysis product ( could be very persistent, the substance has a low potential for bioaccumulation, such that repeated exposures are not expected to cause food-chain effects via accumulation in exposed organisms. Fate: Environmental fate is the determination of which environmental compartment(s) a chemical moves to, the expected residence time in the environmental compartment(s) and removal and degradation processes. Environmental fate is an important factor in determining exposure and thus in determining whether a chemical may present an unreasonable risk. EPA estimated physical/chemical and fate properties of the new chemical substance using data for analogues (acetyl fluorides); of the hydrolysis product (perfluoro acid) using data for analogue(s) (highly fluorinated aliphatic chemicals), and data submitted for the hydrolysis product; and of the hydrolysis product ( using data available for and fluoride ions. In wastewater treatment, the new chemical substance is expected to be removed with an efficiency of 90% to 99% due to rapid hydrolysis, the hydrolysis product (perfluoro acid) is expected to be removed with an efficiency of 0% due to low biodegradability, low sorption, and low stripping, and the hydrolysis product is expected to be removed with an efficiency of 60% due

to sorption. Removal of the hydrolysis products (perfluoro acid and

by biodegradation is negligible. Sorption of the

hydrolysis product (perfluoro acid) to sludge, soil, and sediment is expected

to be low and sorption of the hydrolysis product ( sludge, soil, and sediment is expected to be moderate. Migration of the new chemical substance to groundwater is expected to be negligible due to rapid hydrolysis, migration of the hydrolysis product (perfluoro acid) to groundwater is expected to be rapid due to low sorption to soil and sediment, and migration of the hydrolysis product ( groundwater is expected to be moderate due to moderate sorption to soil and sediment. Due to low estimated vapor pressure and Henry's law constant, the new chemical substance and the hydrolysis product (perfluoro acid) are expected to undergo negligible volatilization to air. Due to low reported vapor pressure, the hydrolysis product ( is expected to undergo negligible volatilization to air. Overall, these estimates indicate that the new chemical substance has low potential to volatilize to air or migrate to groundwater; that the hydrolysis product (perfluoro acid) has low potential to volatilize to air and has high potential to migrate to groundwater; and that the hydrolysis product ( has low potential to volatilize to air and has moderate potential to migrate to groundwater.

Persistence: Persistence is relevant to whether a new chemical substance is likely to present an unreasonable risk because chemicals that are not degraded in the environment at rates that prevent substantial buildup in the environment, and thus increase potential for exposure, may present a risk if the substance presents a hazard to human health or the environment. EPA estimated degradation half-lives of the new chemical substance using data for analogues (acetyl fluorides), of the hydrolysis product (perfluoro acid) using data for analogue(s) (highly fluorinated aliphatic chemicals), and of the hydrolysis product ( using data available for EPA estimated that the new chemical substance's hydrolysis half-life is seconds; that the hydrolysis products' (perfluoro acid aerobic and anaerobic biodegradation half-lives are > 6 months. These estimates indicate that the new chemical substance may have limited persistence in aerobic environments (e.g., surface water) and anaerobic environments (e.g., sediments) due to hydrolysis. Further, these estimates indicate that the hydrolysis products (perfluoro acid and may be very persistent in aerobic environments (e.g., surface water) and anaerobic environments (e.g., sediment).

Bioaccumulation: Bioaccumulation is relevant to whether a new chemical substance is likely to present an unreasonable risk because substances that bioaccumulate in aquatic and/or terrestrial species pose the potential for elevated exposures to humans and other organisms via food chains. EPA estimated the potential for the new chemical substance to bioaccumulate using data for analogues (acetyl fluorides) and of the hydrolysis product to bioaccumulate using data for fluoride ions. EPA estimated that the new chemical substance has low bioaccumulation potential based on rapid hydrolysis, the hydrolysis product (perfluoro acid) has unknown bioaccumulation potential based on bioconcentration or

bioaccumulation data reported for highly fluorinated aliphatic chemicals, and the hydrolysis product ( has low bioaccumulation potential based on fluoride, which is a nutrient. EPA estimated that the new chemical substance could have limited persistence and a low potential for bioaccumulation, such that repeated exposures are not expected to cause food-chain effects via accumulation in exposed organisms. The hydrolysis product (perfluoro acid) is estimated to be very persistent in the environment and its bioaccumulation potential is unknown; therefore, it is not known whether repeated exposures to it will cause food chain effects via accumulation in exposed organisms. Although EPA estimated that the hydrolysis product ( could be very persistent, the substance has a low potential for bioaccumulation, such that repeated exposures are not expected to cause food-chain effects via accumulation in exposed organisms.

### **Physical Chemical Information**

Molecular Weight:			
Wt% < 500:	\$ 39	Wt% < 1000:	
Physical State - Neat:			
Melting Point:		Melting Point (est):	
MP (EPI):			
Vapor Pressure:		Vapor Pressure (est):	
VP (EPI):			
Water Solubility:		Water Solubility (est):	
Water Solubility			<u> </u>
(EPI):			
Henry's Law::			
Log Koc:	NaN	Log Koc (EPI):	
Log Kow:		Log Kow (EPI):	
Log Kow			
Comment:			

#### SAT Concern Level

<b>Ecotox Rating</b>	2		
(1):			
<b>Ecotox Rating</b>			
Comment (1):			
<b>Ecotox Rating</b>			
(2):			
<b>Ecotox Rating</b>			

Comment (2):

Ecotox Route of All releases to water

Exposure:

#### **Ecotox Comments**

Exposure Based
Review (Eco):
Ecotox
Comments:
Exposure Based
Testing:

# **PBT Ratings**


Persistence	Bioaccumulation	Toxicity	Comments	
1	1		PMN	
3	U		Hyd Pdt	
3	1		Hyd Pdt	

# **Eco-Toxicity Comment:**

# **Fate Ratings**

(Overall): Condition	Rating		Rating I	Description		Comment
	Values	1	2	3	4	
Fish BCF:						
Log Fish BCF:						
WWT/POTW	;1;2	Low	Moderate	Strong	V. Strong	PMN;Hyd
Sorption:						Pdt
						Herd
						Hyd Pdt
WWT/POTW	4;4;4	Extensive	Moderate	Low	Negligible	PMN;Hyd
Stripping:	', ', '	Emelisive	Moderate	Lon	riegngiere	Pdt
11 5						
						Hyd

Removal in 9 WWT/POTW (Overall):	0-99;0;60	PMN;Hyd Pd	lt Perfluoro a	ncid; Hyd Pdt		
Condition	Rating	12		escription		Comment
Biodegradation Removal:	Values ;4;4	1 Unknown	2 High	3 Moderate	4 Negligible	PMN;Hyd Pdt acid; Hyd Pdt
Biodegradation Destruction:		Unknown	Complete	Partial	<del></del>	
Aerobic Biodeg Ult:	;4;4	<= Days	Weeks	Months	> Months	PMN;Hyd Pdt acid; Hyd Pdt
Aerobic Biodeg Prim:		<= Days	Weeks	Months	> Months	
Anaerobic Biodeg Ult:	;4;4	<= Days	Weeks	Months	> Months	PMN;Hyd Pdt acid; Hyd Pdt
Anaerobic Biodeg Prim:		<= Days	Weeks	Months	> Months	
Hydrolysis (t1/2 at pH 7,25C) A:		<= Minutes	Hours	Days	>= Months	
Hydrolysis (t1/2 at pH 7,25C) B:		<= Minutes	Hours	Days	>= Months	
Sorption to Soils/Sediments:	;4;3	V. Strong	Strong	Moderate	Low	PMN;Hyd Pdt
Migration to Ground Water:	1;4;3	Negligible	Slow	Moderate	Rapid	acid; Hyd Pdt PMN;Hyd Pdt Perfluoro acid; Hyd Pdt
Photolysis A,		Negligible	Slow	Moderate	Rapid	Tut
Direct: Photolysis B,		Negligible	Slow	Moderate	Rapid	
Indirect: Atmospheric Ox A, OH:		Negligible	Slow	Moderate	Rapid	



## **Ecotoxicity Values**

Test organism	Test Type	Test Endpoint	Predicted	Experimental Comments
Fish	96-h	LC50	>96.9	Predictions are
				based on
				analogue test
				data
Daphnid	48-h	LC50	>100	Predictions are
				based on
				analogue test
				data
	110.2 300	44.22	15/9/2	
Green Algae	96-h	EC50	>106	Predictions are
				based on
				analogue test
				data
Fish	10 <b>-</b>	Chronic Value	1.53	Predictions are
				based on
				analogue test
				data
Daphnid	% <del>=</del>	Chronic Value	5.82	Predictions are
				based on
				analogue test
				data
				\$ <del></del>

Test organism Green Algae	Test Type -	Test Endpoint Chronic Value	Predicted > 106	Experimental Comments  Predictions are based on analogue test data
Avian		Chronic value	> 950	Predictions are based on analogue test data
Comments: K P a	tow = (M MN), dispersib	ts and mean measi	with an unkr effective con	MW Log nown MP (P); S = (M, ncentrations based on 100% trations; hardness <150 mg/L as

#### **Ecotox Factors**

Factors	Most Sensitive Endpoint	Assessment Factor	CoC	Comment
Acute	> 96900	5	> 19380	Fish acute
Aquatic(ppb): Chronic Aquatic(ppb):	1530	10	153	Fish ChV

<b>Factors</b>	Values	Comments
SARs:		
SAR Class:		
TSCA NCC None		
Category?		

#### Recommended

Testing:

**Ecotox Factors** Environmental Hazard: Environmental hazard is relevant to whether a new **Comments:** chemical substance is likely to present unreasonable risk because the significance of the risk is dependent upon both the hazard (or toxicity) of the chemical substance and the extent of exposure to the substance. EPA estimated environmental hazard of this new chemical substance using hazard data on analogous chemicals. Acute toxicity values estimated for fish, aquatic invertebrates, and algae are >96.9 mg/L, >100 mg/L, and >106 mg/L, respectively. Chronic toxicity values estimated for fish, aquatic invertebrates, and algae are 1.53 mg/L, 5.82 mg/L, and >106 mg/L, respectively. These toxicity values indicate that the new chemical substance

is expected to have moderate environmental hazard. Application of assessment factors of 5 and 10 to acute and chronic toxicity values, respectively, results in acute and chronic concentrations of concern of 19.38 mg/L (19,380 ppb) and 0.153 mg/L (153 ppb), respectively.

Environmental Risk: Risks to the environment were not identified due to no releases to water.

# **Comments/Telephone Log**

Artifact	Update/Upload Time	
----------	--------------------	--